## Table 17.11 Mesothelioma

## SEER Incidence Age-Adjusted Rates and Trends<sup>a</sup> By Race/Ethnicity and Sex

SEER Incidence	SEER 17 Areas <sup>b</sup> Rate 2004-2008 Rate per 100,000 persons			SEER 13 Areas <sup>b</sup> Trend 1999-2008 <sup>c</sup> AAPC (%)			SEER 9 Areas <sup>b</sup> Trend 1999-2008 <sup>d</sup> AAPC (%)		
RACE/ETHNICITY	Total	Males	Females	Total	Males	Females	_ Total	Males	Females
All Races	1.1	1.9	0.4	-1.1*	-1.6*	0.1	-1.2*	-1.8*	-0.1
White White Hispanic <sup>e</sup> White Non-Hispanic <sup>e</sup>	1.2 1.0 1.2	2.1 1.7 2.2	0.5 0.5 0.5	-0.8* -0.6 -0.7*	-1.3* -2.8* -1.2*	0.5 - 0.1	-0.8* - -	-1.5* - -	0.8*
Black	0.7	1.2	0.3	-	-	-	-5.8*	-6.7*	-
Asian/Pacific Islander	0.3	0.6	0.2	-	-	-	-	-	_
Amer Ind/Alaska Natf	1.1	-	-	-	-	-	-	-	_
Hispanic <sup>e</sup>	1.0	1.6	0.5	-0.5	-2.7*	-	-	-	-

The AAPC is the Average Annual Percent Change over the time interval. The AAPCs are calculated by the Joinpoint Regression Program Version 3.5, April 2011, National Cancer Institute. Statistic not shown. Rate based on less than 16 cases for the time interval.

Trend based on less than 10 cases for at least one year within the time interval.

Rates are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130). Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).

b The SEER 9 areas are San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.

The SEER 13 areas comprise the SEER 9 areas plus San Jose-Monterey, Los Angeles, the Alaska Native Registry and Rural Georgia.

The SEER 17 areas comprise the SEER 13 areas plus California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey.

The 1999-2008 AAPC estimates are based on a Joinpoint analysis with up to 3 Joinpoints over diagnosis years 1992-2008.

d The 1999-2008 AAPC estimates are based on a Joinpoint analysis with up to 5 Joinpoints over diagnosis years 1975-2008.

Hispanic and Non-Hispanic are not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics and Non-Hispanics are based on NHIA and exclude cases from the Alaska Native Registry

Incidence data for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.

The APC is significantly different from zero (p<.05).